Garden Safe® Brand

Slug & Snail Bait

Cebo para babosas y caracoles

KILLS SLUGS & SNAILS

Remains Effective After Rain or Sprinkling!

17-12774

Active Ingredient:	By weight
Iron phosphate	1.0%
Other Ingredients	99.0%
Total	100.0%

SCHULTZ®

KEEP OUT OF REACH OF CHILDREN

CAUTION See back booklet for additional precautionary statements.

Net Wt 3.0 lb (1.3 kg)

MANTÉNGASE FUERA DEL ALCANCE DE LOS NIÑOS

PRECAUCIÓN Consulte el panel en la parte posterior para conocer las precauciones adicionales.

COVERS UP TO 3,000 sq ft

Kills Slugs & Snails!

- *For Household Home Garden Use*
- *Can Be Used in Vegetable Gardens*

Garden Safe® Brand Slug & Snail Bait is a unique blend of an iron phosphate active ingredient, originating from soil, with slug and snail bait additives. It is used as an ingredient in fertilizers. The bait which is not ingested by snails and slugs will degrade and become a part of the soil.

The bait is attractive to slugs and snails and lures them from their hiding places and plants. Ingestion, even in small amounts, will cause them to cease feeding. This physiological effect of the bait gives immediate protection to the plants even though the slugs and snails may remain in the area. After eating the bait, the slugs and snails cease feeding, become less mobile and begin to die within three to six days. Dead slugs and snails may not be visible as they often crawl away to secluded places to die. Plant protection will be observed in the decrease in plant damage.

This product is effective against a wide variety of slugs and snails and will give protection to home lawns, gardens, greenhouses, outdoor ornamentals, vegetable gardens, fruits, berries, citrus and crop plants. The bait can be scattered on the lawn or on the soil around any vegetable plants, flowers or fruit trees or bushes to be protected.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

How To Apply

Scatter the Garden Safe Brand Slug & Snail Bait granules on the soil around or near the plants to be protected. For broadcast application, use standard broadcast spreaders. For row application, use standard granular spreaders. Apply bait evenly at approximately ½-1 lb per 1,000 square feet, or for smaller areas, 0.15 oz or 1 level teaspoon per square yard, and reapply as the bait is consumed or at least every two weeks. Do not place in piles. If the ground is dry, wet it before applying bait. The product works best when soil is moist but with little or no standing water. Reapply as the bait is consumed or at least every two weeks. Apply more heavily if the infestation is severe, if the area is heavily watered or after long periods of heavy rain. See specific directions for different plant types and for inside greenhouses.

When To Apply

Evening is the best time to apply the bait, as slugs and snails travel and feed mostly by night or early morning.

Where To Apply

Treat all likely areas of infestation, especially around the perimeter of garden plots because these pests travel into plant areas from daytime refuges. They favor damp places around vegetable plants such as beans, tomatoes, lettuce, cabbage, celery and squash. Other favorite areas are flower gardens, rockeries, hedges, dichondra lawns, citrus groves, ivy patches and other ground cover where they obtain shelter by day.

Vegetables • The bait can be used to protect any vegetables from slug and snail damage, including (but not limited to): Artichokes, asparagus, beans, beets, blackeyed peas, broccoli, Brussels sprouts, cabbage, cantaloupe, carrots, cauliflower, corn, cucumbers, eggplants, garlic, lettuce, onions, peas, peppers, potatoes, radishes, rutabagas, spinach, squash, Swiss chard, tomatoes and turnips. Scatter the bait around the perimeter of the vegetable plot at approximately ½-1 lb per 1,000 square feet, or for smaller areas, 0.15 oz or 1 level teaspoon per square yard, to provide a protective "barrier" for slugs and snails entering the garden plot. If slugs or snails are inside the rows, then scatter the bait on the soil around the base of the plants and between the rows.

Fruits Including Citrus • The bait can be used to protect fruits from slugs and snails, including (but not limited to): Apples, apricots, avocados, cherries, citrus, grapes, melons, peaches, pears and plums. For seedlings, spread the bait around the base of the stem. Apply at 0.15 oz or 1 level teaspoon per square yard, in a 6-inch circular band around the base of the plants to be protected. For older trees, spread the bait around the base of the tree to intercept slugs and snails traveling to the trunk. Apply the bait at approximately ½-1 lb per 1,000 square feet for orchards using standard fertilizer granular spreaders.

Questions or Comments? Call 1-800-257-3379 or visit our website at **www.gardensafe.com**

WARRANTY • Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller. Seller makes no other warranties, either expressed or implied.

The Garden Safe® Promise • Satisfaction guaranteed or your money back!
If you apply Garden Safe® Brand Slug & Snail Bait as directed and are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address at left. We will refund the purchase price you paid.

0 72845 00453 6

PROOF OF PURCHASE

EPO

MADE IN THE USA

Distributed by Schultz Company
PO Box 4406, Bridgeton, MO 63044-0406

EPA Reg. No. 67702-3-39609
EPA Est. Nos. 39609-MO-1 (BM), 9688-MO-1 (U)
Circled letter is first letter of lot number.
17-12917 © 2008

Made with Ferramol® Slug and Snail Bait, a registered trademark of W. Neudorff GmbH KG. Manufactured under a license of W. Neudorff GmbH KG, Germany.

◄ Open label here for directions for use and other precautions.

Berries • The bait can be used to protect berries from slugs and snails, including (but not limited to): Blackberries, blueberries, boysenberries, loganberries, raspberries and strawberries. Spread the bait around the perimeter of the plot to intercept slugs and snails migrating toward the berries. Use a rate of approximately ½-1 lb per 1,000 square feet, or for smaller areas, 0.15 oz or 1 level teaspoon per square yard, and scatter by hand or with granular spreaders. If slugs and snails are already in the plots, then carefully spread bait between the furrows near the base of the plants. For small plots, treat around the base of the plants to be protected. Do not spread over the entire area but apply selectively.

Outdoor Ornamentals • Scatter bait in a 6-inch circular band around the base of the plants to be protected at 0.15 oz or 1 level teaspoon per square yard. If plants are next to a grassy area, spread the bait between the ornamentals and the grass. Slugs and snails traveling to the plants will encounter the bait before reaching the plant. In these situations, scatter the bait around the perimeter of the plot at approximately 1 lb per 1,000 square feet to intercept snails and slugs traveling to the plot.

Greenhouses • Where slugs or snails are a problem in greenhouses, scatter the bait in the plant pots of plants being damaged or around pots on greenhouse benches. Apply about ½ teaspoon per 9-inch pot.

Lawns • The bait can be used to protect lawns. When slugs or snails are detected, scatter the bait at a rate of approximately ½-1 lb per 1,000 square feet, or for smaller areas, 0.15 oz or 1 level teaspoon per square yard. Scatter by hand or with a granular spreader where the slugs or snails are observed.

MODO DE EMPLEO

Se considera una violación a la ley federal usar este producto de una manera que no sea la indicada en la etiqueta.

Cómo aplicar

Esparza los gránulos de cebo para babosas y caracoles en el suelo, alrededor o cerca de las plantas que se van a proteger. Para aplicaciones al voleo, use difusores al voleo estándar. Para aplicación en filas, use difusores granulares estándar. Aplique el cebo de manera uniforme a una dosis de aproximadamente ½-1 lb. por cada 1,000 pies cuadrados, o en áreas más pequeñas, 0.15 oz. o 1 cucharadita rasa por yarda cuadrada, y vuelva a aplicar cuando se consuma el cebo por lo menos cada dos semanas. No lo apile. Si la tierra está seca, mójela antes de aplicar el cebo. Este producto actúa mejor cuando el suelo está húmedo pero sin agua estancada, o con poca agua. Aplique nuevamente cuando se consuma el cebo o por lo menos cada dos semanas. Aplique más cantidad si la infestación es grave, si el área se riega mucho o después de períodos prolongados de lluvia fuerte. Consulte las instrucciones específicas para diferentes tipos de plantas y para invernaderos interiores.

Cuándo aplicar

El mejor momento para aplicar el cebo es durante la tarde, ya que las babosas y los caracoles transitan y comen más en la noche o temprano por la mañana.

Dónde aplicar

Trate todas las áreas donde haya probabilidades de infestación, especialmente alrededor del perímetro de parcelas de jardín porque estas plagas ingresan en las áreas plantadas desde los refugios diurnos. Prefieren los lugares húmedos alrededor de hortalizas como frijoles, tomate, lechuga, coles, apio y calabaza. Otros lugares favoritos son los jardines de flores, rocallas, setos vivos, céspedes dichondra, huertas de cítricos, manchas de hiedra y otros tapices vegetales donde se refugian de día.

Hortalizas • El cebo se puede usar para proteger del daño ocasionado por babosas y caracoles a hortalizas que incluyen, entre otras: alcachofas, espárragos, frijoles, betabel, frijoles de carita, brócoli, coles de Bruselas, col, melones, zanahorias, coliflor, maíz, pepinos, berenjena, ajo, lechuga, cebollas, chícharos, pimientos, papas, rábanos, rutabaga, espinaca, calabaza, acelga, tomates y nabos. Esparza el cebo alrededor del perímetro de la parcela de hortalizas en la proporción de ½-1 lb. por cada 1,000 pies cuadrados aproximadamente, o en áreas más pequeñas, 0.15 oz. o 1 cucharadita rasa por yarda cuadrada, para proporcionar una "barrera" de protección contra babosas y caracoles que entren a la parcela de jardín. Si las babosas o caracoles están dentro de las filas, esparza el cebo en la tierra alrededor de la base de las plantas y entre las filas.

Frutas, incluidos cítricos • El cebo se puede usar para proteger del daño ocasionado por babosas y caracoles, a frutas que incluyen, entre otras: manzanas, chabacanos, aguacates, cerezas, cítricos, uvas, melones, duraznos, peras y ciruelas. Para plántulas, esparza el cebo alrededor de la base del tallo. Aplique a la dosis de 0.15 oz. o 1 cucharadita rasa por yarda cuadrada, en una franja circular de 6 pulgadas alrededor de la base de las plantas que se van a proteger. En árboles más viejos, esparza el cebo alrededor de la base del árbol para detener a las babosas y los caracoles que transiten hacia el tronco. Aplique el cebo a una dosis aproximada de ½-1 lb. por cada 1,000 pies cuadrados en huertos, utilizando difusores estándar para fertilizantes granulados.

Bayas • El cebo se puede usar para proteger del daño ocasionado por babosas y caracoles, a bayas que incluyen, entre otras: zarzamora, blueberries, moras de Boysen, moras de Logan, frambuesa y fresas. Esparza el cebo alrededor del perímetro de la parcela para detener a babosas y caracoles que transiten hacia las bayas. Use una dosis aproximada de ½-1 lb. por cada 1,000 pies cuadrados, o en áreas más pequeñas, 0.15 oz. o 1 cucharadita rasa por yarda cuadrada, y esparza a mano o con difusores granulares. Si las babosas y los caracoles ya están en las parcelas, esparza cuidadosamente el cebo entre los surcos cerca de la base de las plantas. En parcelas pequeñas, trate alrededor de la base de las plantas para que queden protegidas. No esparza sobre toda el área sino que aplique selectivamente.

Plantas ornamentales de exterior • Esparza el cebo en una franja circular de 6 pulgadas alrededor de la base de las plantas que se desea proteger a la dosis de 0.15 oz. o 1 cucharadita rasa por yarda cuadrada. Si las plantas están cerca de una superficie de pasto, esparza el cebo entre las plantas ornamentales y el pasto. Las babosas y los caracoles que transitan hacia las plantas encontrarán el cebo antes de llegar a la planta. En estas situaciones, esparza el cebo alrededor del perímetro de la parcela a la dosis aproximada de 1 lb. por cada 1,000 pies cuadrados para detener a las babosas y los caracoles que transiten hacia la parcela.

Invernaderos • Donde las babosas y los caracoles sean un problema en invernaderos, esparza el cebo en las macetas de las plantas que sufrieron daño o alrededor de las macetas en las mesas del invernadero. Aplique aproximadamente ½ cucharadita para una maceta de 9 pulgadas.

Céspedes • El cebo se puede usar para proteger céspedes. Cuando se detecten babosas o caracoles, esparza el cebo a una dosis aproximada de ½-1 lb. por cada 1,000 pies cuadrados, o en áreas más pequeñas, 0.15 oz. o 1 cucharadita rasa por yarda cuadrada. Esparza a mano o con un difusor granular donde vea babosas o caracoles.

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

PESTICIDE STORAGE: Store this product in its original container and keep in a secure storage area out of reach of children and domestic animals.

CONTAINER DISPOSAL: **If empty:** Do not reuse this container. Place in trash or offer for recycling if available. **If partly filled:** Call your local solid waste agency for disposal instructions. Never place unused product down any indoor or outdoor drain.

PRECAUTIONARY STATEMENTS

Hazards to Humans and Domestic Animals: **Caution.** Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling.

First Aid

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **Hotline Number:** Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact the National Pesticide Telecommunications Network hotline at 1-800-858-7378 for emergency medical treatment information.

Environmental Hazards • For terrestrial uses. Do not apply directly to water. Do not contaminate water when disposing of equipment washwaters or rinsate.

Questions or Comments? Call 1-800-257-3379 or visit our website at **www.gardensafe.com**

ALMACENAMIENTO Y ELIMINACIÓN

No contamine el agua ni los alimentos para personas o animales por medio del almacenamiento o la eliminación.

ALMACENAMIENTO DEL PESTICIDA: Guarde este producto en su envase original, en un lugar seguro fuera del alcance de los niños y los animales domésticos.

ELIMINACIÓN DEL ENVASE: **Si está vacío:** No vuelva a usar este envase. Tírelo en la basura u ofrézcalo para reciclar, si existe esa opción. **Si está parcialmente vacío:** Llame a su servicio de eliminación de desechos más cercano para obtener instrucciones de eliminación. Nunca tire el producto restante por un drenaje interno ni externo.

PRECAUCIONES

Riesgos para los seres humanos y animales domésticos: **Precaución.** Causa irritación moderada a los ojos. Evite el contacto con los ojos o la ropa. Lávese bien con agua y jabón después de manipular el producto.

Primeros Auxilios

Si entra en contacto con los ojos: Mantenga los ojos abiertos y enjuague con agua lentamente, con cuidado, durante 15-20 minutos. Si hay lentes de contacto, retírelos después de los primeros 5 minutos, luego continúe enjuagando los ojos. Llame a un centro de control de envenenamientos o a un médico para recibir instrucciones de tratamiento. **Número de la línea directa:** Cuando llame a un centro de control de envenenamientos o a un médico, o intente obtener tratamiento, tenga a la mano el envase o la etiqueta del producto. También puede llamar a la National Pesticide Telecommunications Network al 1-800-858-7378 para obtener información sobre tratamiento médico de urgencia.

Riesgos para el medio ambiente • Para uso terrestre. No aplique directamente al agua. No contamine el agua cuando elimine las aguas de lavado del equipo ni los residuos del enjuague.

¿Preguntas o comentarios? Llame al 1-800-257-3379.

WARRANTY • Seller warrants that this product conforms to the chemical description on this label and is reasonably fit for purposes stated on this label only when used in accordance with directions under normal use conditions. This warranty does not extend to use of this product contrary to label directions, or under abnormal use conditions, or under conditions not reasonably foreseeable to seller. Seller makes no other warranties, either expressed or implied.

The Garden Safe® Promise • Satisfaction guaranteed or your money back!
If you apply Garden Safe® Brand Slug & Snail Bait as directed and are not completely satisfied, send proof of purchase from this package and the original cash register receipt to the address at left. We will refund the purchase price you paid.

MADE IN THE USA

Distributed by Schultz Company
PO Box 4406, Bridgeton, MO 63044-0406

EPA Reg. No. 67702-3-39609
EPA Est. Nos. 39609-MO-1 (BM), 9688-MO-1 (U)
Circled letter is first letter of lot number.
17-12917 © 2008

Made with Ferramol® Slug and Snail Bait, a registered trademark of W. Neudorff GmbH KG. Manufactured under a license of W. Neudorff GmbH KG, Germany.